



## Product Datasheet

<b>Product Name</b>	Interleukin-17E Human Recombinant
<b>Cata No</b>	CB501317
<b>Source</b>	<i>Escherichia Coli.</i>
<b>Synonyms</b>	IL-25, IL-17E, IL17E, IL25, Interleukin-25, Interleukin-17E.

### Description

IL-25 also called IL-17E cytokine has a sequence similarity with IL17. IL-17E induces NF-kappaB activation, and stimulates the production of IL-8. IL17E and IL17B are ligands for the cytokine receptor IL17BR. IL-25 is a proinflammatory cytokine favoring Th2-type immune response. The upregulation of costimulation-induced IL-17E receptors and release of cytokines and chemokines from IL-17E treated costimulated Th cells are differentially regulated by intracellular JNK, p38 MAPK and NF-kappaB activity. Blocking interleukin-25 prevents airway hyperresponsiveness, a critical feature of clinical asthma. IL25 produced by innate effector eosinophils and basophils increase the allergic inflammation by enhancing the maintenance and functions of TSLP-DC activated adaptive Th2 memory cells. Over expression of IL-25 up-regulates gene expression of Th2 cytokines and induces growth retardation, jaundice, and multiorgan inflammation in a transgenic mouse model. IL-25 contributes to the induction and maintenance of eosinophilic inflammation by acting on lung fibroblasts which supports the fact that IL-17E is an important factor in asthma pathophysiology. IL-17E operates by amplifying TH2 cell-mediated allergic airway inflammation but doesn't induce allergic inflammation in vivo. Interleukin-17E Human Recombinant produced in E.Coli is a homodimeric, non-glycosylated polypeptide chain containing 146 amino acids and

having a molecular mass of 33.7 kDa. The IL-25 Human is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

### Purity

Greater than 95.0% as determined by:  
(a) Analysis by RP-HPLC.  
(b) Analysis by SDS-PAGE.

### Formulation

Lyophilized from a concentrated (1mg/ml) solution containing no additives.

### Reconstitution

It is recommended to reconstitute the lyophilized Interleukin Human IL25 in sterile 10mM HCL not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized Interleukin Human IL17E although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Human IL-25 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

**Please prevent freeze-thaw cycles.**

### Sequence

MYSHWPSCCP SKGQDTSEEL LRWSTVPVPP

**\* For Non-Clinical Research Use Only \***



# California Bioscience

83103 Avenue 48, Ste.1B #204  
Coachella, CA 92236 USA  
Phone : +1.6268339877  
Email : [info@cali-bio.com](mailto:info@cali-bio.com)

LEPARPNRHP ESCRASEDGP LNSRAISPWR  
YELDRDLNRL PQDLYHARCL CPHCVSLQTG

SHMDPRGNSE LLYHN **Product Data Sheet**  
HKGYCLERRL YRVSLACVCV RPRVMG

**\* For Non-Clinical Research Use Only \***